

On the topological decomposition of the hypersurfaces in projective toric manifolds

Wei WANG

School of Mathematical Sciences, Fudan University
Shanghai, P. R. China, 200433

December, 2011

Abstract

In this paper, we want to discuss the topology of the non-singular hypersurface Y^n with complex dimension n in a projective toric manifold X^{n+1} . When n is odd, our main results are a decomposition of $Y^n \cong Y' \sharp s(S^n \times S^n)$ as a connected sum of s copies of $S^n \times S^n$ with a differential manifold Y' such that $b_n(Y') = 0$ or 2 . When n is even and the degree of Y in X is big enough, we find that Y also admits such a decomposition $Y' \sharp s(S^n \times S^n)$, where Y' satisfy $b_n(Y') - |\text{sign}(Y')| = b_n(X) \pm \text{sign}(H^n(X))$, where $\text{sign}(H^n(X))$ is the signature of a certain bilinear form defined on $H^n(X, \mathbb{Z})$.

1 Introduction

1.1 Projective toric manifold and its hypersurfaces

Definition 1.1. *A toric variety is a normal algebraic variety X containing the algebraic torus $(\mathbb{C}^*)^n$ as a Zariski open subset in such a way that the normal action $(\mathbb{C}^*)^n$ on itself extends to an action on X .*

*In this paper, we call X a **projective toric manifold** if X is a compact, smooth toric variety that admits a holomorphic embedding into a certain $\mathbb{C}P^N$.*

The algebraic topology of projective toric manifold has been fully studied by many people and many results can be found in these two classical books [4],[5]. In this paper, what we need are the following two propositions ([5], page 56,101,102).

Proposition 1.2. *Let X be a projective toric manifold, then X is simply connected and the odd dimension homology groups of X vanish, i.e. $H_{\text{odd}}(X, \mathbb{Z}) = 0$.*

Proposition 1.3. *$H_*(X, \mathbb{Z})$ can be generated by the projective toric submanifolds of X , i.e. there exist smooth toric submanifolds $\{X_i\}$ with $x_i = [X_i] \in H_*(X, \mathbb{Z})$ such that the homomorphism*

$$\sum \mathbb{Z}x_i \longrightarrow H_*(X, \mathbb{Z})$$

¹MSC(2000): 57 R 19, 57 R 65

²Keywords: Removing handles, Connected sum, Hypersurfaces, Toric manifolds

is surjective.

Then we introduce the hypersurface of a projective toric manifold. Let X be a projective toric manifold. For any holomorphic embedding $X \hookrightarrow \mathbb{C}P^N$, let F_1 be a hyperplane of $\mathbb{C}P^N$, we get a subvariety $i : Y = F_1 \cap X \hookrightarrow X$ of X and Y is called a **hypersurface** of X . By Bertini's theorem, for a generic hyperplane F_1 in $\mathbb{C}P^N$, Y is smooth.

Given such a hypersurface Y of $X \hookrightarrow \mathbb{C}P^N$, we can also construct the smooth hypersurface $i_d : Y_d \hookrightarrow X$ of X with $(i_d)_*[Y_d] = d(i_*[Y])$, $0 < d \in \mathbb{Z}$. Indeed, we can take $Y_d := F_d \cap X$, where F_d is a generic hypersurface of $\mathbb{C}P^N$ with degree d and it is well-known that Y_d is also a smooth hypersurface of X .

In this paper, all the hypersurfaces we consider are smooth and when we say a hypersurface Y_d , it always means Y_d is a smooth hypersurface.

Similar to the degree of a hypersurface in $\mathbb{C}P^n$, we can define the degree of a (smooth) hypersurface in X . Let α_Y be the element of $H^2(X, \mathbb{Z})$ such that $\alpha_Y \cap [X] = i_*[Y]$. We define the **degree** of a hypersurface Y in X by

$$\deg Y := \langle \alpha_Y^{n+1}, [X] \rangle$$

For the hypersurface Y_d , we have relation $d\alpha_Y = \alpha_{Y_d}$ and we have $\deg Y_d = d^{n+1} \deg Y$.

1.2 Main results

Let X^{n+1} be a projective toric manifold with complex dimension $n+1$, $n > 2$. Let $i : Y \hookrightarrow X$ be the hypersurface of X with complex dimension n and $i_d : Y_d \hookrightarrow X$ be the hypersurface with $(i_d)_*[Y_d] = d(i_*[Y])$, $0 < d \in \mathbb{Z}$. In this paper, we want to discuss the topological decomposition of the hypersurface Y_d . Our main results are:

Theorem 1.4. *When n is odd, for any integer $d > 0$, we have decomposition:*

$$Y_d \cong Y'_d \# s_d(S^n \times S^n)$$

where the n -th Betti number $b_n(Y'_d) = 0$ or 2.

Theorem 1.5. *When n is even, for sufficiently big $d \gg 0$, we have decomposition:*

$$Y_d \cong Y'_d \# s_d(S^n \times S^n)$$

with $s_d = \frac{b_n(Y_d) - b_n(X) - |\text{sign}(Y_d) - \text{sign}(H^n(X))|}{2}$, here $\text{sign}(Y_d)$ is the classical signature of Y_d and $\text{sign}(H^n(X))$ is the signature of the bilinear form defined by:

$$\begin{aligned} H^n(X) \otimes H^n(X) &\longrightarrow \mathbb{Z} \\ (x, y) &\mapsto \langle x \cup y \cup \alpha_{Y_d}, [X] \rangle \end{aligned}$$

Furthermore, we have limit estimate:

$$0 < \lim_{d \rightarrow +\infty} \frac{2s_d}{\deg Y_d} = \lim_{d \rightarrow +\infty} \frac{2s_d}{d^{n+1} \deg Y} = 1 - 2^{n+1}(2^{n+1} - 1) \frac{B_{\frac{n+2}{2}}}{(n+1)!} < 1$$

here $B_{\frac{n+2}{2}}$ is the $\frac{n+2}{2}$ -th Bernoulli number.

For Y'_d , we have relations $b_n(Y'_d) = b_n(Y_d) - 2s_d$ and $\text{sign}(Y_d) = \text{sign}(Y'_d)$. By the limit estimates in proposition 4.4, we know $|\text{sign}(Y_d)| = |\text{sign}(Y'_d)|$ tends to $+\infty$ as $d \rightarrow +\infty$. From theorem 1.5, we can deduce that

Corollary 1.6. *When n is even and d is big enough:*

$$b_n(Y'_d) - |\text{sign}(Y'_d)| = b_n(X) \pm |\text{sign}(H^n(X))|$$

Remark 1.7. Let F be a nonsingular algebraic hypersurface in complex projective space, in [8], Kulkarni and Wood proved that there is a differentiable connected sum decomposition

$$F = M \sharp k(S^n \times S^n)$$

where $b_n(M) = 0$ or 2 for n odd, and $b_n(M) - |\text{sign}(M)| = b_n(\mathbb{C}P^n) \pm \text{sign}(H^n(\mathbb{C}P^{n+1})) = 0$ or 2 for n even.

Our theorem is a generalization of their theorem to the case of hypersurfaces in projective toric manifolds.

2 Basic idea of removing handles

2.1 Geometric point of view

Choose a point $(x, y) \in S^n \times S^n$ and there are two embedded spheres: $S_1 := S^n \times \{y\}$, $S_2 := \{x\} \times S^n \hookrightarrow S^n \times S^n$ with properties:

- (1). S_1 intersects S_2 transversally at one point (x, y) .
- (2). The normal bundles of S_1, S_2 in $S^n \times S^n$ are trivial.
- (3). Denote $\eta_1 := S_1 \times D^n \subset S^n \times S^n$ and $\eta_2 := S_2 \times D^n \subset S^n \times S^n$ by the closure of their normal bundles, we see $\eta_1 \cup \eta_2$ is a manifold with boundary S^{2n-1} and

$$S^n \times S^n = (\eta_1 \cup \eta_2) \cup_{S^{2n-1}} D^{2n}$$

Conversely, let M^{2n} be a smooth manifold and S_1, S_2 be two embedded n -spheres of M^{2n} with:

- (1). S_1 intersects S_2 transversally at one point.
- (2). The normal bundles of S_1 and S_2 are trivial.

We denote $\xi_1 := S_1 \times D^n$, $\xi_2 := S_2 \times D^n$ by the closure of their normal bundles. Observe that $\eta_1 \cup \eta_2 \cong \xi_1 \cup \xi_2$ and we get:

$$M \cong (M - \xi_1^\circ \cup \xi_2^\circ) \cup_{S^{2n-1}} (\eta_1 \cup \eta_2) \cong M' \sharp S^n \times S^n$$

where $M' = (M - \xi_1^\circ \cup \xi_2^\circ) \cup_{S^{2n-1}} D^{2n}$. This is the basic idea of removing handles from a $2n$ -manifold ([12]). Next we want to realize this idea by algebraic topology.

2.2 Homological point of view

From the point of view of homology, let M^{2n} be a simply connected smooth closed manifold of dimension $2n$, $n > 2$ and $h : \pi_n(M) \rightarrow H_n(M, \mathbb{Z})$ be the Hurewicz map. For every

$\alpha, \beta \in h(\pi_n(M)) \subset H_n(M, \mathbb{Z})$ with intersection number $\alpha \cdot \beta = 1$, by Whitney's embedding theory and Whitney's trick ([10], p142), there are two embedding n -spheres $f_\alpha, f_\beta : S^n \hookrightarrow M^{2n}$ with:

- (1). The homology elements α and β are represented by f_α, f_β , i.e. $(f_\alpha)_*[S^n] = \alpha, (f_\beta)_*[S^n] = \beta$
- (2). The spheres $f_\alpha(S^n)$ and $f_\beta(S^n)$ intersect transversally at only one point.

Following the geometric idea of removing handles, the next question is how to determine the normal bundles. In general, the normal bundles of $f_\alpha(S^n), f_\beta(S^n)$ are not easy to determine.

In this paper, the situation seems relatively simpler: let $K \subset h(\pi_n(M))$ be a free Abel group such that each element $\alpha \in K$ can be represented by an embedded n -sphere $f_\alpha : S^n \hookrightarrow M$ with **stable trivial** normal bundle.

When n is even, for the embedding f_α representing $\alpha \in K$, the normal bundle of f_α is just determined by the self-intersection number $\alpha \cdot \alpha$ of α . Indeed, $\alpha \cdot \alpha = 0$ if and only if the normal bundle of f_α is trivial. So, if we could find a free subgroup $\oplus_{i=1}^s (\mathbb{Z}\alpha_i \oplus \beta_i)$ of K with intersection matrix $\oplus_{i=1}^s \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$, topologically, M admits a decomposition:

$$M \cong M' \sharp s(S^n \times S^n)$$

When n is odd, the intersection number $\alpha \cdot \alpha$ is always zero and can not determine the normal bundle of f_α , we need two techniques.

Technique 1: find a quadratic function $\psi : K \longrightarrow \mathbb{Z}_2$ with:

- (1). $\psi(\alpha + \beta) = \psi(\alpha) + \psi(\beta) + (\alpha \cdot \beta)_2$, where $(\alpha \cdot \beta)_2 \in \mathbb{Z}_2$ is the mod 2 class of the intersection number $\alpha \cdot \beta \in \mathbb{Z}$, which is also the definition of the quadratic function over \mathbb{Z} .
- (2). $\psi(\alpha) = 0$ if and only if α can be represented by an embedded n -sphere f_α with trivial normal bundle.

For any free subgroup $\oplus_{i=0}^s (\mathbb{Z}\alpha'_i \oplus \beta'_i)$ of K with intersection matrix $\oplus_{i=0}^s \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$, by the standard results of the quadratic function ([15], p172), we can find a new basis $\{\alpha_i, \beta_i\}$ of this subgroup such that the intersection matrix of $\mathbb{Z}\alpha_0 \oplus \mathbb{Z}\beta_0 \oplus \{\oplus_{i=1}^s (\mathbb{Z}\alpha_i \oplus \beta_i)\}$ is still $\oplus_{i=0}^s \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ and $\psi(\alpha_i) = \psi(\beta_i) = 0, i \neq 0, \psi(\alpha_0) = \psi(\beta_0) = 0$ or 1 . In this case, although we can not determine the value of $\psi(\alpha_0)$, at least, we have decomposition:

$$M \cong M' \sharp s(S^n \times S^n)$$

In general, the quadratic function ψ is not always exist on K and we need the second technique.

Technique 2: find an embedded n -sphere $g : S^n \hookrightarrow M$ with:

- (1). $g_*[S^n] = 0 \in H_n(M, \mathbb{Z})$.
- (2). The normal bundle η_g of $g : S^n \hookrightarrow M$ is isomorphic to the tangent bundle TS^n of S^n .

If we could find such an embedding g , for every element $\alpha \in K$ which is represented by an embedding f_α , if the stable trivial normal bundle η_{f_α} is not trivial, by Wall's technique ([15], p167), there exists a new embedding f'_α with normal bundle $\eta_{f'_\alpha}$ such that:

- (1). $f'_\alpha = f_\alpha + g \in \pi_n(M)$
- (2). $F(\eta_{f'_\alpha}) = F(\eta_{f_\alpha}) + F(\eta_g)$ where F is the isomorphism:

$$\{n \text{ dimensional stable trivial vector bundles over } S^n\} \longleftrightarrow \text{Ker}(\pi_{n-1}(SO(n)) \rightarrow \pi_{n-1}(SO))$$

It is known that $\text{Ker}(\pi_{n-1}(SO(n)) \rightarrow \pi_{n-1}(SO)) = 0$, $n = 1, 3, 7$ and $\text{Ker}(\pi_{n-1}(SO(n)) \rightarrow \pi_{n-1}(SO)) = \mathbb{Z}_2$ with generator $F(TS^n)$, n odd $\neq 1, 3, 7$. ([2], p88).

In this case, modifying by this embedding g , we can make every element $\alpha \in K$ represented by an embedding with trivial normal bundle. So, for any free subgroup $\oplus_{i=1}^s (\mathbb{Z}\alpha_i \oplus \beta_i)$ of K with intersection matrix $\oplus_{i=1}^s \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$, M admits a decomposition:

$$M \cong M' \#_s (S^n \times S^n)$$

These are the basic tools to removing (n dimensional) handles from a $2n$ -manifold. In the next two sections, we will apply these tools to prove our main theorems.

3 Odd case

3.1 Wu class, quadratic function, and Kervaire invariant

Given a smooth manifold $(M^n, \partial M)$, the Steenrod operator $Sq = \sum_{i=0} Sq^i : H^*(M, \partial M, \mathbb{Z}_2) \rightarrow H^*(M, \partial M, \mathbb{Z}_2)$ determines a linear form on $H^*(M, \partial M, \mathbb{Z}_2)$:

$$H^*(M, \partial M) \longrightarrow \mathbb{Z}_2$$

$$x \mapsto \langle Sq(x), [M] \rangle$$

where $[M] \in H_n(M, \partial M, \mathbb{Z}_2)$ is the fundamental class of the Poincaré pair $(M, \partial M)$. Since the cup product induces the isomorphism $H^*(M, \mathbb{Z}_2) \cong \text{Hom}(H^*(M, \partial M, \mathbb{Z}_2), \mathbb{Z}_2)$, there exists a unique element $v(M) = 1 + v_1(M) + v_2(M) + \dots \in H^*(M, \mathbb{Z}_2)$ such that for each $x \in H^*(M, \partial M, \mathbb{Z}_2)$:

$$\langle v(M) \cup x, [M] \rangle = \langle Sq(x), [M] \rangle, \quad \langle v_i(M) \cup x, [M] \rangle = \langle Sq^i x, [M] \rangle$$

Definition 3.1. $v(M) = \sum_{i=0} v_i(M)$ is called the Wu class of M .

By the definition, we see $v_i(M) = 0 \iff Sq^i : H^{n-i}(M, \partial M, \mathbb{Z}_2) \rightarrow \mathbb{Z}_2$ is zero.

In his paper [3], Browder gave a geometric definition of Kervaire invariant, which is equivalent to his original definition of Kervaire invariant in [1]. This geometric definition is very close to the original definition of Kervaire, which is defined by the Arf invariant of a certain quadratic function, (cf [7]). First, it is known in [3] that:

Proposition 3.2. For any $x \in H_n(M^{2n}, \mathbb{Z}_2)$, we can find an embedded $N^n \subset M^{2n}$ with $[N] = x$.

Proposition 3.3. If $M^{2n} \times \mathbb{R}^q \subset W^{2n+q}$, W connected and $y \in H_{n+1}(W, M, \mathbb{Z}_2)$, we can find $N \subset M$ representing $\partial y \in H_n(M, \mathbb{Z}_2)$, i.e $[N] = \partial y$ with $N = \partial V$, here $i : V \subset W \times [0, 1]$ is a connected submanifold with $i_*[V] = y$, where $[V] \in H_{n+1}(V, \partial V, \mathbb{Z}_2)$ is the fundamental class. Furthermore, V meets $W \times 0$ transversally in $N \subset M$.

We see, in this case, the normal bundle of N in $W \times 0$ has a normal q -frame ($N \times \mathbb{R}^q \subset M \times \mathbb{R}^q \subset W$). The obstruction to extending this frame to a normal q -frame on $V \subset W \times [0, 1]$ lies in $H^{i+1}(V, N, \pi_i(V_{n+q,q}))$ and $\pi_i(V_{n+q,q}) = 0, q < n, \pi_n(V_{n+q,q}) = \mathbb{Z}_2$. We find the last and only one obstruction $\sigma \in H^{n+1}(V, N, \pi_n(V_{n+q,q})) = \mathbb{Z}_2$.

Definition 3.4. For the element $x = \partial y \in H_n(M, \mathbb{Z}_2)$, where $\partial : H_{n+1}(W, M, \mathbb{Z}_2) \rightarrow H_n(M, \mathbb{Z}_2)$ and $y \in H_{n+1}(W, M, \mathbb{Z}_2)$, we define: $\psi(x) = \langle \sigma, [V] \rangle$, which is denoted briefly by σ for convenience.

We see this definition seems not intrinsic, it depends on the choice of N and V . We should put some condition to make ψ well-defined. Browder proved ([3]):

Proposition 3.5. The obstruction to extend a q -frame defines a quadratic form:

$$\psi : \text{Ker}(H_n(M, \mathbb{Z}_2) \rightarrow H_n(W, \mathbb{Z}_2)) \rightarrow \mathbb{Z}_2$$

if and only if $v_{n+1}(W) = 0$.

Proposition 3.6. For the embedding $\phi(S^n) \in M^{2n, n}$ odd, if $\phi(S^n)$ is nullhomotopic in W , then $\psi([\phi(S^n)]) = 0$ if and only if the normal bundle of $\phi(S^n)$ is trivial.

Definition 3.7. If $\text{Ker}(H_n(M, \mathbb{Z}_2) \rightarrow H_n(W, \mathbb{Z}_2))$ is non-singular under the intersection pair, we define the Kervaire invariant k by its Arf invariant of the quadratic form ψ .

3.2 Proof of the odd case I

Let X^{n+1} be a projective toric manifold with complex dimension $n > 2$, odd, and $i : Y^n \hookrightarrow X^{n+1}$ be a hypersurface of X^{n+1} .

Lemma 3.8. $H_n(Y, \mathbb{Z})$ is spherical and every element $\alpha \in H_n(Y, \mathbb{Z})$ can be represented by an embedding $f_\alpha : S^n \hookrightarrow Y$ such that the normal bundle η_{f_α} of f_α is stable trivial.

Proof. First, by Lefschetz's hyperplane section theorem and Proposition 1.2., we know (X, Y) is n -connected and $H_n(X) = 0$. We have exact sequence:

$$\begin{array}{ccccc} H_{n+1}(X, Y) & \longrightarrow & H_n(Y) & \longrightarrow & H_n(X) = 0 \\ \uparrow \cong & & \uparrow h_Y & & \uparrow \\ \pi_{n+1}(X, Y) & \longrightarrow & \pi_n(Y) & \xrightarrow{\pi_n(i)} & \pi_n(X) \end{array}$$

From this diagram, we observe that $h_Y : \pi_n(Y) \rightarrow H_n(Y)$ is surjective and for every element $\alpha \in H_n(Y)$, by the Whitney embedding theorem, we can choose an embedding $f_\alpha : S^n \hookrightarrow Y$ to represent α such that $i \circ f_\alpha$ is nullhomotopic in X , i.e. $\pi_n(i)([f_\alpha]) = 0$.

Second, we want to show the normal bundle η_{f_α} is stable trivial. We have bundle identity:

$$TX|S^n = (i \circ f_\alpha)^*TX = TS^n \oplus \eta_{f_\alpha} \oplus \eta_Y^X|S^n$$

here η_Y^X is the normal bundle of $i : Y \rightarrow X$. Since η_Y^X is a complex line bundle, it is known that $\eta_Y^X \cong i^*L_Y$, where L_Y is a complex line bundle over X with Euler class $e(L_Y) \cap [X] = i_*[Y]$.

Since $i \circ f_\alpha$ is nullhomotopic, the bundle identity becomes:

$$\epsilon^{2n+2} = (i \circ f_\alpha)^* TX = TS^n \oplus \eta_{f_\alpha} \oplus (i \circ f_\alpha)^* L_Y = \epsilon^{n+1} \oplus \eta_{f_\alpha}$$

here ϵ is the trivial real 1-bundle. \square

Proof of the odd case I: For the complex line bundle L_Y in the above lemma, consider $W = D(-L_Y)$, where $-L_Y$ is the stable inverse bundle of L_Y , i.e. $L_Y \oplus -L_Y$ is trivial, and $D(-L_Y)$ is the disk bundle of $-L_Y$. Then for the embedding: $Y \hookrightarrow X \hookrightarrow W$, we see the normal bundle of Y in W is trivial and we get $Y \times \mathbb{R}^q \subset W$ for some $q > 0$.

Observe that $\text{Ker}(H_n(Y, \mathbb{Z}_2) \rightarrow H_n(W, \mathbb{Z}_2)) = H_n(Y, \mathbb{Z}_2)$ and by proposition 3.5, if the Wu class $v_{n+1}(W) = 0$, there is a quadratic function $\psi' : H_n(Y, \mathbb{Z}_2) \rightarrow \mathbb{Z}_2$ and we also obtain a quadratic function on $H_n(Y, \mathbb{Z})$:

$$\psi : H_n(Y, \mathbb{Z}) \longrightarrow H_n(Y, \mathbb{Z}_2) \xrightarrow{\psi'} \mathbb{Z}_2$$

Furthermore, by proposition 3.6, we know $\psi(\alpha) = 0$ if and only if the normal bundle η_{f_α} is trivial.

Since $H_n(Y, \mathbb{Z})$ is unimodular, by technique 2 in subsection 2.2, $H_n(Y, \mathbb{Z}) \cong \mathbb{Z}\alpha_0 \oplus \mathbb{Z}\beta_0 \oplus \bigoplus_{i=1}^s (\mathbb{Z}\alpha_i \oplus \beta_i)$, $s = \frac{b_n(Y)-2}{2}$ with intersection matrix $\bigoplus_{i=0}^s \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ and $\psi(\alpha_i) = \psi(\beta_i) = 0, i \neq 0$. Topologically, we get decomposition:

$$Y \cong Y' \sharp s(S^n \times S^n)$$

where $b_n(Y') = 2$. If the Kervaire (Arf) invariant k of ψ or ψ' vanishes, we can make $b_n(Y') = 0$.

So we finish the proof of the odd case when $v_{n+1}(W) = 0$. When $v_{n+1}(W) \neq 0$, the quadratic function ψ is not necessary well-defined and we will use technique 2 to deal with it in next subsection.

3.3 Proof of the odd case II

In his paper [3], Browder proved:

Theorem 3.9 (Browder). *Suppose $M^{2n} \times \mathbb{R}^q \subset W$, $n \neq 1, 3$ or 7 , W is 1-connected. (W, M) is n -connected and suppose $v_{n+1}(W) \neq 0$. Then there exists an embedded $S^n \subset M^{2n}$ and $U^{n+1} \subset M^{2n} \times \mathbb{R}^{q+1}$ with $\partial U = S^n$ such that the normal bundle ξ to S^n in M^{2n} is nontrivial, but $\xi \oplus \epsilon^1$ is trivial, where ϵ^1 is the trivial one dimensional real vector bundle. Hence S^n is homologically trivial (mod 2) with nontrivial normal bundle.*

Remark 3.10. It seems we can use this theorem to find the embedding sphere in technique 2. But the shortage is: the embedding sphere S^n is only mod 2 trivial. We want to add some condition to make it work in integral homology.

Theorem 3.11. *Under the same hypothesis of Browder's theorem above, if we further assume that $H_{n+1}(W, \mathbb{Z}_2)$ is generated by the element $\{x_i\}$ such that each x_i can be represented by an oriented closed manifold N_i , i.e $[N_i] = x_i$. Then there exist an embedded $S^n \subset M^{2n}$ such that $[S^n] = 0 \in H_n(M, \mathbb{Z})$ and the normal bundle ξ to S^n in M^{2n} is non-trivial but stable trivial.*

Proof. We follow Browder's proof (Step 2 to Step 7 is almost unchanged):

Step 1: Since $v_{n+1}(W) \neq 0$, we know $Sq^{n+1} : H^{n+q-1}(W, \partial W, \mathbb{Z}_2) \rightarrow \mathbb{Z}_2$ is not zero. By assumption, $\exists N_i$ such that $Sq^{n+1}y_i \neq 0$, where $y_i \cap [W] = [N_i]$.

Step 2: For convenience, we denote N_i by N and y_i by y . Let $N_0 = N - \text{int}D^{n+1}$, we see $\partial N_0 = S^{n+1}$ and N_0 is homotopic to an n -complex. Since (W, M) is n -connected, $\exists f : N_0 \rightarrow M$ such that the diagram is commutative up to homotopy:

$$\begin{array}{ccc} N_0 & \xrightarrow{f} & M \\ \downarrow & & \downarrow \\ N & \longrightarrow & W \end{array}$$

Step 3: Let $g = f|_{\partial N_0} : S^n \rightarrow M$, we see $g_*[S^n] = 0 \in H_n(M, \mathbb{Z})$. Since M is 1-connected, by Whitney's theorem, we can make g homotopic to an embedding and we still denote it by g . Since g is nullhomotopic in W , then the normal bundle of $g(S^n)$ is stable trivial. We wish to show that the normal bundle of this sphere is not trivial.

Step 4: We make the map $f : N_0 \rightarrow M \times \mathbb{R}^q \times [-1, 0]$ homotopic to an embedding $g_0 : N_0 \hookrightarrow M \times \mathbb{R}^q \times [-1, 0]$ such that $g_0|_{\partial N_0} = g$. And we extend $g : S^n \hookrightarrow M$ to $\tilde{g} : D^{n+1} \subset W \times [0, 1]$ which meets $W \times 0$ transversally in $g(S^n)$. Then we get an embedding $g_1 : N_0 \cup_{S^n} D^{n+1} \cong N \hookrightarrow W \times [-1, 1]$, which is isotopic to the origin $N \subset W$.

Step 5: $M \times \mathbb{R}^q \subset W \times 0$ define a q -frame of the normal bundle of $g(S^n) \subset M \subset W \times 0$. We know the obstruction $\sigma \in \pi_{n+1}(V_{n+1,q})$ to extend this q -frame to D^{n+1} is zero if and only if the normal bundle of $g(S^n)$ is trivial.

Step 6: Now assume the normal bundle of $g(S^n)$ is trivial and we get a q -frame on D^{n+1} :

$$D^{n+1} \times D^n \times \mathbb{R}^q \subset W \times [0, 1]$$

such that $D^{n+1} \times 0 \times 0 = \tilde{g}(D)$ and $S^n \times D^n \times 0$ is the normal bundle of $g(S^n)$. Let $V = M \times [-1, 0] \cup_{S^n \times D^n} D^{n+1} \times D^n$, then we get $V \times \mathbb{R}^q \subset W \times [-1, 1]$, $g_1(N) \subset \text{int}V \times \mathbb{R}^q$.

Step 7: Let $Y = W \times [-1, 1]/\partial(W \times [-1, 1])$ we get:

$$Y \xrightarrow{a} \Sigma^q V / \partial V \xrightarrow{b} T(\eta_N \oplus \epsilon)$$

where η_N is the normal bundle of N in W . Let U be the mod 2 Thom class of $\eta_N \oplus \epsilon$, we get: $(ba)^*U = \Sigma x \in H^{n+q}(Y, \mathbb{Z}_2)$ and $(Sq^{n+1}(x))[W] = Sq^{n+1}(\Sigma x)[Y] \neq 0$. Also, $(Sq^{n+1}(b^*U))(\Sigma^q[V]) \neq 0$ and $Sq^{n+1}(\Sigma^{-q}(b^*U))[V] \neq 0$. On the other hand, $Sq^{n+1}(\Sigma^{-q}(b^*U)) = 0$ since $\Sigma^{-q}(b^*U) = 0 \in H^n(V, \partial V, \mathbb{Z}_2)$. \square

Proof of the odd case II: When $v_{n+1}(W) \neq 0$, in our case $Y \times \mathbb{R}^q \subset W = D(-L)$, by proposition 1.3, we see $H_*(X, \mathbb{Z})$ is generated by the toric submanifolds which are certainly oriented and $W = D(-L_Y)$ is the disk bundle over X , whose homology group is also generated by these toric submanifolds. Then all the conditions of theorem 3.11 are satisfied. Thus, there exists an embedding sphere $g : S^n \hookrightarrow Y$ such that $g_*[S^n] = 0$ and the normal bundle $\eta_g \cong TS^n$.

By the technique 2 in section 2 and lemma 3.8, we have topological decomposition:

$$Y \cong Y' \sharp s(S^n \times S^n)$$

where $b_n(Y') = 0$. Then we finish the proof of theorem 1.4.

4 Even case

4.1 Intersection form and signature

Let X^{n+1} be a projective toric manifold with complex dimension $n + 1$, $n > 2$, even, and $i : Y \hookrightarrow X$ be a hypersurface of X . Since n is even, the n -th homology group $H_n(Y, \mathbb{Z})$ admits a unimodular symmetric intersection form:

$$H_n(Y, \mathbb{Z}) \otimes H_n(Y, \mathbb{Z}) \longrightarrow \mathbb{Z}$$

Since (X, Y) is n -connected and $H_{\text{odd}}(X, \mathbb{Z}) = 0$, like the odd case, we have

$$\begin{array}{ccccccc} 0 & \longrightarrow & H_{n+1}(X, Y) & \longrightarrow & H_n(Y) & \xrightarrow{i_*} & H_n(X) \longrightarrow 0 \\ & & \uparrow \cong & & \uparrow h_Y & & \uparrow \\ & & \pi_{n+1}(X, Y) & \longrightarrow & \pi_n(Y) & \xrightarrow{\pi_n(i)} & \pi_n(X) \end{array}$$

The **vanishing cycles** $\text{Ker}(i_*) \subset H_n(Y, \mathbb{Z})$ is what we mainly concerned, because:

Lemma 4.1. *Each element $\alpha \in \text{Ker}(i_*)$ can be represented by an embedding $f_\alpha : S^n \hookrightarrow Y$ such that $f_\alpha[S^n] = \alpha$ and the normal bundle η_{f_α} of f_α is stable trivial.*

Proof. Since $\pi_n(X, Y) \cong H_n(X, Y, \mathbb{Z}) \cong \text{Ker}(i_*)$, we see for each element $\alpha \in \text{Ker}(i_*)$, there exists an embedding f_α representing α and $\pi_n(i)(f_\alpha) = 0 \in \pi_n(X)$.

The proof of the stable triviality of the normal bundle η_{f_α} is similar to the proof in lemma 3.8. \square

When we restrict the intersection form of $H_n(Y, \mathbb{Z})$ on $\text{Ker}(i_*)$, we get:

Proposition 4.2. *The intersection form on $\text{Ker}(i_*)$ is of type even, i.e. for any $\alpha \in \text{Ker}(i_*)$, $\alpha \cdot \alpha$ is even.*

Proof. For any $\alpha \in \text{Ker}(i_*)$, by lemma 4.1., we can use an embedding f_α to represent it. It is known that $\alpha \cdot \alpha = \langle e(\eta_{f_\alpha}), [S^n] \rangle$, where $e(\eta_{f_\alpha})$ is the Euler class of the normal bundle η_{f_α} . Furthermore, $\langle e(\eta_{f_\alpha}), [S^n] \rangle$ is even if and only if the n -th Stiefel-Whitney class $w_n(\eta_{f_\alpha})$ of η_{f_α} is zero and this is just proved in lemma 4.1. \square

The intersection pair on $H_n(Y, \mathbb{Z})$ is equivalent to the cup product on $H^n(Y, \mathbb{Z})$

$$H^n(Y, \mathbb{Z}) \otimes H^n(Y, \mathbb{Z}) \longrightarrow \mathbb{Z}$$

$$(\alpha, \beta) \mapsto \langle \alpha \cup \beta, [Y] \rangle$$

through the Poincaré duality $PD : H^n(Y, \mathbb{Z}) \longrightarrow H_n(Y, \mathbb{Z})$ and we also have exact sequence:

$$0 \longrightarrow H^n(X, \mathbb{Z}) \xrightarrow{i^*} H^n(Y, \mathbb{Z}) \longrightarrow H^{n+1}(X, Y) \longrightarrow 0$$

We see the intersection form $(\text{Ker}(i_*), \cdot)$ is equivalent to $(PD^{-1}(\text{Ker}(i_*)), \cup)$ and the reason why we use the language of cohomolgy instead of homology is:

Lemma 4.3. $PD^{-1}(Ker(i_*)) = (i^*H^n(X))^\perp$

Proof. For any $\alpha \in Ker(i_*)$ and $\beta \in H^n(X, \mathbb{Z})$, we have:

$$\langle PD^{-1}(\alpha) \cup i^*\beta, [Y] \rangle = \langle i^*\beta, \alpha \rangle = \langle \beta, i_*\alpha \rangle = 0$$

we get $PD^{-1}(Ker(i_*)) \subset (i^*H^n(X))^\perp$.

On the other hand, for any $PD^{-1}(\gamma) \in (i^*H^n(X))^\perp$, we see

$$\langle PD^{-1}(\gamma) \cup i^*H^n(X, \mathbb{Z}), [Y] \rangle = \langle i^*H^n(X, \mathbb{Z}), \gamma \rangle = \langle H^n(X, \mathbb{Z}), i_*\gamma \rangle = 0$$

Since $H_n(X, \mathbb{Z})$ and $H^n(X, \mathbb{Z})$ are free Abel groups, we get $i_*\gamma = 0$. \square

Next, we want to discuss some limit estimates about the n -th Betti number and the signature of the pair $(H^n(Y_d, \mathbb{Z}), \cup)$. Recall that $i_d : Y_d \hookrightarrow X^{n+1}$ is the hypersurface of the toric manifold X with $(i_d)_*[Y_d] = d(i_*[Y])$ and $deg Y_d = \langle \alpha_{Y_d}^{n+1}, [X] \rangle = d^{n+1}deg Y$, where $\alpha_{Y_d} \cap [X] = (i_d)_*[Y_d]$. We have proposition:

Proposition 4.4. *We have limits:*

$$\begin{aligned} \lim_{d \rightarrow +\infty} \frac{b_n(Y_d)}{deg Y_d} &= \lim_{d \rightarrow +\infty} \frac{b_n(Y_d)}{d^{n+1}deg Y} = 1 \\ 0 &< \lim_{d \rightarrow +\infty} \frac{|sign(Y_d)|}{b_n(Y_d)} = 2^{n+2}(2^{n+2} - 1) \frac{B_{\frac{n+2}{2}}}{(n+1)!} < 1 \end{aligned}$$

Proof. For the first limit, we know the Euler number $\chi(Y_d)$ of Y_d equals $b_n(Y_d) + 2 \sum_{j=1}^{n-1} (-1)^j b_j(X)$ and

$$\begin{aligned} \chi(Y_d) &= \langle c_n(Y_d), [Y_d] \rangle = \langle \frac{c(TX)}{1 + d\alpha_Y}, [X] \rangle \\ &= d^{n+1} \langle \alpha_Y^{n+1}, [X] \rangle + O(d^n) \end{aligned}$$

here $c(TX)$ and $c_n(Y_d)$ are the Chern classes. We have:

$$\lim_{d \rightarrow +\infty} \frac{\chi(Y_d)}{d^{n+1}} = \lim_{d \rightarrow +\infty} \frac{b_n(Y_d)}{d^{n+1}} = deg Y$$

and we get:

$$\lim_{d \rightarrow +\infty} \frac{b_n(Y_d)}{deg Y_d} = 1$$

For the second limit, we have identity:

$$sign(Y_d) = \langle \tanh(d\alpha_Y) L(X), [X] \rangle$$

where $L(X) = L_1(X) + L_2(X) + \dots$ is the L -class of X and $\tanh(d\alpha_Y) = \sum_{j=1}^{+\infty} (-1)^{j-1} 2^{2j} (2^{2j} - 1) \frac{B_j}{(2j)!} (d\alpha_Y)^{2j-1}$. Observe that:

$$sign(Y_d) = (-1)^{\frac{n}{2}} 2^{n+2} (2^{n+2} - 1) \frac{B_{\frac{n+2}{2}}}{(n+2)!} d^{n+1} deg Y + O(d^n)$$

we have limit:

$$\lim_{d \rightarrow +\infty} \frac{|sign(Y_d)|}{b_n(Y_d)} = 2^{n+2}(2^{n+2} - 1) \frac{B_{\frac{n+2}{2}}}{(n+1)!}$$

Furthermore, when $j > 1$, we see:

$$1 + \frac{1}{2^{2j}} + \frac{1}{3^{2j}} + \dots = \frac{B_j(2\pi)^{2j}}{2(2j)!} < \frac{\pi^2}{6}$$

$$\frac{B_j 2^{2j}(2^{2j} - 1)}{(2j)!} < \frac{\pi^2}{3} \frac{2^{2j}(2^{2j} - 1)}{(2\pi)^{2j}} < \frac{\pi^2}{3} \frac{4^j}{\pi^{2j}} < 1$$

□

Corollary 4.5. $\lim_{d \rightarrow +\infty} b_n(Y_d) = +\infty$ and $\lim_{d \rightarrow +\infty} sign(Y_d) = +\infty$. When d is big enough, $(H_n(Y_d, \mathbb{Z}), \cdot)$ is indefinite.

4.2 Proof of the even case

Let $(H, <, >)$ be a unimodular symmetric bilinear form over \mathbb{Z} and F be a nonzero subgroup of H such that H/F is free and the map $F \rightarrow Hom(F, \mathbb{Z})$ induced by $<, >$ is injective. Denote $E = F^\perp := \{x \in H \mid <x, F> = 0\}$, we have:

Theorem 4.6. *If $rank H \geq \max\{4rank F, 2rank F + 5\}$, then E admits an orthogonal decomposition:*

$$(E, <, >) \cong (A, <, >) \oplus (\oplus_{i=1}^s (\mathbb{Z}x_i \oplus \mathbb{Z}y_i, <, >))$$

where the intersection matrix of $\mathbb{Z}x_i \oplus \mathbb{Z}y_i$ is $\begin{pmatrix} 0 & 1 \\ 1 & c_i \end{pmatrix}$, $c_i = 0$ or 1 . For $(A, <, >)$, there are two possibilities:

- (1). $(A, <, >)$ is definite and $rank A \geq \max\{3rank F, rank F + 5\}$
- (2). $rank A < \max\{3rank F, rank F + 5\}$

We'll prove the even case first and the proof of this theorem will be given in the next subsection.

Proof of the even case: Step 1: For the bilinear symmetric space $(H^n(Y_d, \mathbb{Z}), \cup)$, we know $PD^{-1}(Ker((i_d)_*)) = (i^*H^n(X))^{\perp}$. We want to show the injectivity of the map $H^n(X, \mathbb{Z}) \rightarrow Hom(H^n(X, \mathbb{Z}), \mathbb{Z})$ induced by the cup product in $H^n(Y_d, \mathbb{Z})$.

Since Y_d is the hypersurface of X , the hard Lefschetz theorem ([11]) tell us that the cohomology element α_{Y_d} representing Y_d induces an injective map:

$$\cup \alpha_{Y_d} : H^n(X, \mathbb{Z}) \rightarrow H^{n+2}(X, \mathbb{Z})$$

For $i^*H^n(X, \mathbb{Z}) \subset H^n(Y_d, \mathbb{Z})$, we have diagram:

$$\begin{array}{ccccc} H^n(X, \mathbb{Z}) & \xrightarrow{\cup \alpha_{Y_d}} & H^{n+2}(X, \mathbb{Z}) & \xrightarrow{\cong} & Hom(H^n(X, \mathbb{Z}), \mathbb{Z}) \\ i_d^* \downarrow & & & & \uparrow \\ i_d^* H^n(X, \mathbb{Z}) & \longrightarrow & H^n(Y_d, \mathbb{Z}) & \xrightarrow{\cong} & Hom(H^n(Y_d, \mathbb{Z}), \mathbb{Z}) \end{array}$$

indeed, for any $x, y \in H^n(X, \mathbb{Z})$, $x(y) = \langle i^*x \cup i^*y, [Y_d] \rangle = \langle x \cup y \cup \alpha_{Y_d}, [X] \rangle = (x \cup \alpha_{Y_d})(y)$.

Furthermore, we see the restriction of $(H^n(Y_d, \mathbb{Z}), \cup)$ to $H^n(X, \mathbb{Z})$ is just the bilinear form defined in theorem 1.5.

Thus we get a pair $(H_d, \cup) = (H^n(Y_d, \mathbb{Z}), \cup)$ with a free subgroup $F := i^*H^n(X, \mathbb{Z})$ such that

- (1). $F^\perp = PD^{-1}(Ker(i_d)_*) = E_d$ with even type (proposition 4.2, lemma 4.3).
- (2). If d is big enough, $rank H_d > Max\{4rank F, 2rank F + 5\}$ (proposition 4.4)

Step 2: By the algebraic decomposition theorem 4.6,

$$(Ker(i_d)_*, \cdot) \cong (E_d, \cup) \cong A_d \oplus (\oplus_{i=1}^{s_d} (\mathbb{Z}x_i \oplus \mathbb{Z}y_i, \langle, \rangle))$$

where the intersection matrix of $\mathbb{Z}x_i \oplus \mathbb{Z}y_i$ is $\begin{pmatrix} 0 & 1 \\ 1 & c_i \end{pmatrix}$, $c_i = 0$ or 1

By proposition 4.2, $Ker((i_d)_*)$ is of type even, c_i must be zero. Since $\lim_{d \rightarrow \infty} |sign H^n(Y_d, \mathbb{Z})| = +\infty$, when d is big enough, the possibility (2) of theorem 4.6 can not happen, and A_d is definite.

Step 3: By the process of removing handles of the even case in section 2, we see

$$Ker(i_d)_* = A_d \oplus \oplus_{i=1}^{s_d} (\mathbb{Z}x_i \oplus \mathbb{Z}y_i)$$

where the intersection matrix of $\oplus_{i=1}^{s_d} (\mathbb{Z}x_i \oplus \mathbb{Z}y_i)$ is $\oplus_{s_d} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ and we get:

$$Y_d \cong Y'_d \#_{s_d} (S^n \times S^n)$$

Since A_d is definite and $sign(\mathbb{Z}x_i \oplus \mathbb{Z}y_i) = 0$, we get identiy $2s_d = rank(Ker(i_d)_*) - |sign(Ker(i_d)_*)|$. Also, since $Ker(i_d)_* = (H^n(X, \mathbb{Z}))^\perp \subset H^n(Y_d, \mathbb{Z})$ and the restriction of $(H^n(Y_d, \mathbb{Z}), \cup)$ to $H^n(X, \mathbb{Z})$ is just the bilinear form defined in theorem 1.5, we get

$$2s_d = b_n(Y_d) - b_n(X) - |sign(Y_d) - sign(H^n(X))|$$

For the limit estimate, we have:

$$\lim_{d \rightarrow +\infty} \frac{2s_d}{deg Y_d} = \lim_{d \rightarrow +\infty} \frac{2s_d}{b_n(Y_d)} = 1 - \lim_{d \rightarrow +\infty} \frac{|sign(Y_d)|}{b_n(Y_d)}$$

4.3 Proof of the algebraic decomposition theorem

In order to prove theorem 4.6., we need some lemmas.

Lemma 4.7. Assume E satisfy $rank E \geq 3rank F$, we can choose a basis $\{f_1, f_2, \dots, f_{r+h}\}$ of $Hom(E, \mathbb{Z})$ such that $\oplus_{i=1}^r \mathbb{Z}f_i \rightarrow Hom(E, \mathbb{Z})/E$ is surjective, $r \leq rank F$, and $\oplus_{j=1}^h \mathbb{Z}f_{r+j} \subset E \subset Hom(E, \mathbb{Z})$, $h \geq 2r$.

Proof. First, since H/F is free, we have:

$$\begin{array}{ccccccc} 0 & \longrightarrow & E & \xrightarrow{\langle, \rangle} & Hom(H, \mathbb{Z}) & \longrightarrow & Hom(F, \mathbb{Z}) \longrightarrow 0 \\ & & \parallel & & \downarrow & & \downarrow \\ 0 & \longrightarrow & E & \xrightarrow{\langle, \rangle} & Hom(E, \mathbb{Z}) & \longrightarrow & Hom(E, \mathbb{Z})/E \longrightarrow 0 \end{array}$$

Since $H/E \cong \text{Hom}(H, \mathbb{Z})/E \cong \text{Hom}(F, \mathbb{Z})$, we see $\text{Hom}(H, \mathbb{Z}) \longrightarrow \text{Hom}(E, \mathbb{Z})$ is surjective and $\text{Hom}(F, \mathbb{Z}) \longrightarrow \text{Hom}(E, \mathbb{Z})/E$ is also surjective.

Second, $\text{rank Hom}(F, \mathbb{Z}) = \text{rank } F$, we can choose $\text{rank } F$ elements $\{g_1, g_2, \dots\}$ of $\text{Hom}(E, \mathbb{Z})$ such that $\mathbb{Z}g_1 + \mathbb{Z}g_2 + \dots + \mathbb{Z}g_{\text{rank } F} \longrightarrow \text{Hom}(E, \mathbb{Z})/E$ is surjective. Then there is a subgroup $\mathbb{Z}g_1 + \mathbb{Z}g_2 + \dots + \mathbb{Z}g_{\text{rank } F} \subset N \subset \text{Hom}(E, \mathbb{Z})$ with $\text{Hom}(E, \mathbb{Z})/N$ free and $r = \text{rank } N = \text{rank}(\mathbb{Z}g_1 + \mathbb{Z}g_2 + \dots + \mathbb{Z}g_{\text{rank } F}) \leq \text{rank } F$.

Third, Since $\text{Hom}(E, \mathbb{Z})/N$ is free, let $\{f_1, \dots, f_r\}$ be a basis of N and extend it to a basis $\{f_1, \dots, f_r, f'_{r+1}, \dots, f'_{r+h}\}$ of $\text{Hom}(E, \mathbb{Z})$. We know $N \longrightarrow \text{Hom}(E, \mathbb{Z})/E$ is surjective, then for any f'_{r+i} , we can find $f_{r+i} = f'_{r+i} - \sum_{j=1}^r a_{ij} f_j$ with $[\sum_{j=1}^r a_{ij} f_j] = [f'_{r+i}] \in \text{Hom}(E, \mathbb{Z})/E$, we see $f_{r+i} \in E$.

Thus we obtain a basis $\{f_1 \dots f_r, f_{r+1}, \dots, f_{r+h}\}$ of $\text{Hom}(E, \mathbb{Z})$ such that $\{f_{r+1}, \dots, f_{r+h}\} \subset E \subset \text{Hom}(E, \mathbb{Z})$. \square

Lemma 4.8. Assume E is indefinite and $\text{rank } E \geq \text{Max}\{3\text{rank } F, \text{rank } F + 5\}$, we can find two elements $x, y \in E$ with $\langle x, x \rangle = 0$, $\langle x, y \rangle = 1$, $\langle y, y \rangle = 0$ or 1 .

Proof. First, by the lemma above, we have a basis $\{f_1 \dots f_r, f_{r+1}, \dots, f_{r+h}\}$ of $\text{Hom}(E, \mathbb{Z})$ with $\mathbb{Z}f_{r+1} \oplus \mathbb{Z}f_{r+2} \oplus \dots \oplus \mathbb{Z}f_{r+h} \subset E \subset \text{Hom}(E, \mathbb{Z})$, $h > \text{max}\{2r, 5\}$. Let $\{f_1^*, \dots, f_{r+s}^*\}$ be the dual of the basis $\{f_1, \dots, f_{r+h}\}$ in $\text{Hom}(E, \mathbb{Z})^* = E$. Define:

$$D = \mathbb{Z}f_{r+1}^* \oplus \dots \oplus \mathbb{Z}f_{r+h}^* \subset E = \text{Hom}(E, \mathbb{Z})^*$$

Second, when D is indefinite, since $\text{rank } D \geq 5$, it is known from Meyer's theorem that there exists an indivisible element $x \in D$ such that $\langle x, x \rangle = 0$ ([8], p255). Then we can also choose an element $y' \in \mathbb{Z}f_{r+1} \oplus \dots \oplus \mathbb{Z}f_{r+h} \subset E$ such that $\langle x, y' \rangle = 1$. Let $y = y' - [\frac{\langle y', y' \rangle}{2}]x'$, we have $\langle x, x \rangle = 0$, $\langle x, y \rangle = 1$, $\langle y, y \rangle = 0$ or 1 .

Third, when D happens to be definite, define $D' := \mathbb{Z}(f_{r+1}^* - c_1 f_1^*) \oplus \mathbb{Z}(f_{r+2}^* - c_2 f_1^*) \oplus \mathbb{Z}(f_{r+3}^* - c_3 f_2^*) \oplus \mathbb{Z}(f_{r+4}^* - c_4 f_2^*) \oplus \dots \oplus \mathbb{Z}(f_{3r-1}^* - c_{2r-1} f_r^*) \oplus \mathbb{Z}(f_{3r}^* - c_{2r} f_r^*) \oplus \dots \oplus \mathbb{Z}(f_{r+h}^* - c_h f_r^*)$.

If we can choose proper $\{c_i \in \mathbb{Z}\}$ to make D' indefinite, we can still find an indivisible element $x \in D'$ such that $\langle x, x \rangle = 0$, and we can also find $y \in \oplus_{j=1}^h \mathbb{Z}f_{r+j} \in E$ such that $\langle x, y \rangle = 1$. So, all we need to do is to prove the next lemma. \square

Lemma 4.9. Following lemma 4.8, suppose D is definite, we can choose proper $\{c_i \in \mathbb{Z}\}$ to make D' indefinite.

Proof. Assume D is positive definite under \langle, \rangle . Consider the real space $\overline{E} := E \otimes \mathbb{R}$, $\overline{D} := D \otimes \mathbb{R}$, $\overline{D'} := D' \otimes \mathbb{R}$ and let $\{f_1^*, \dots, f_{r+h}^*\}$ be the Euclidean orthogonal standard basis of \overline{E} . Define:

$$F : \overline{E} \longrightarrow \mathbb{R} \\ \sum a_i f_i^* \mapsto \sum a_i a_j \langle f_i^*, f_j^* \rangle$$

Observe that F is just the extension of the map $: E \rightarrow \mathbb{Z}, x \mapsto \langle x, x \rangle$ to \overline{E} .

Note that E is indefinite and \mathbb{Q} -unimodular under \langle, \rangle , since by assumption H is unimodular and F is \mathbb{Q} -unimodular. Then we can find a $v \in \overline{E}$ such that $F(v) < 0$ and the Euclidean norm

$|v| = 1$, i.e. $v = \sum_{i=1}^r a_i f_i^* + \sum_{j=1}^h b_j f_{r+j}^*$, $\sum a_i^2 + \sum b_j^2 = 1$. Since D is definite, we see $F(\overline{D} - \{0\}) > 0$ and $(a_1, \dots, a_r) \neq (0, \dots, 0)$.

In the Euclidean norm with orthogonal standard basis $\{f_i^*\}$, we have decomposition $\overline{E} = \overline{D'} \oplus (\overline{D'})^\perp$. By calculation, we see $(\overline{D'})^\perp$ has a standard orthogonal basis:

$$(\overline{D'})^\perp = \text{span}\left\{ \frac{f_1^* + c_1 f_{r+1}^* + c_2 f_{r+2}^*}{\sqrt{1 + c_1^2 + c_2^2}}, \frac{f_2^* + c_3 f_{r+3}^* + c_4 f_{r+4}^*}{\sqrt{1 + c_3^2 + c_4^2}}, \dots, \frac{f_{r-1}^* + c_{2r-3} f_{3r-3}^* + c_{2r-2} f_{3r-2}^*}{\sqrt{1 + c_{2r-3}^2 + c_{2r-2}^2}}, \frac{f_r^* + c_{2r-1} f_{3r-1}^* + c_{2r} f_{3r}^* + \dots + c_h f_{r+h}^*}{\sqrt{1 + c_{2r-1}^2 + \dots + c_h^2}} \right\}$$

For convenience, denote this basis by $\{g_1, g_2, \dots, g_r\}$.

For the vector $v = \sum_{i=1}^r a_i f_i^* + \sum_{j=1}^h b_j f_{r+j}^*$, we can decompose $v = v_1 + v_2$ such that $v_1 \in \overline{D'}$ and $v_2 \in (\overline{D'})^\perp$. By calculation,

$$v_2 = \frac{a_1 + c_1 b_{r+1} + c_2 b_{r+2}}{\sqrt{1 + c_1^2 + c_2^2}} g_1 + \dots + \frac{a_{r-1} + c_{2r-3} b_{3r-3} + c_{2r-2} b_{3r-2}}{\sqrt{1 + c_{2r-3}^2 + c_{2r-2}^2}} g_{r-1} \\ + \frac{a_r + c_{2r-1} b_{3r-1} + c_{2r} b_{3r} + \dots + c_h b_{r+h}}{\sqrt{1 + c_{2r-1}^2 + \dots + c_h^2}} g_r$$

Since $\sum a_i^2 + \sum b_j^2 = 1$, for $\forall \epsilon > 0$, we can choose proper $c_i \in \mathbb{Z}$ ([8], p256) such that

$$\left| \frac{a_1 + c_1 b_{r+1} + c_2 b_{r+2}}{\sqrt{1 + c_1^2 + c_2^2}} \right| < \frac{\epsilon}{r}, \dots, \left| \frac{a_{r-1} + c_{2r-3} b_{3r-3} + c_{2r-2} b_{3r-2}}{\sqrt{1 + c_{2r-3}^2 + c_{2r-2}^2}} \right| < \frac{\epsilon}{r} \\ \left| \frac{a_r + c_{2r-1} b_{3r-1} + c_{2r} b_{3r} + \dots + c_h b_{r+h}}{\sqrt{1 + c_{2r-1}^2 + \dots + c_h^2}} \right| < \frac{\epsilon}{r}$$

The function F is continuous and $F(v) < 0$, if the Euclidean norm of $v_2 = v - v_1$ is small enough, then the element $v_1 \in \overline{D'}$ satisfy $F(v_1) < 0$. Furthermore, D' is not negative definite, since D is positive and $\text{rank} D' = \text{rank} D$, $2\text{rank} D > \text{rank} E = \text{rank} D + \text{rank} F$, thus we see D' is indefinite. □

Proof of theorem 4.6.: We use induction on $\text{rank} H$, since $\text{rank} H \geq \text{Max}\{4\text{rank} F, 2\text{rank} F + 5\}$, we get $\text{rank} E \geq \text{Max}\{3\text{rank} F, \text{rank} F + 5\}$. If E is definite, we've done. If E is indefinite, then by the lemmas we've just proved, there exist two elements $x, y \in E$ such that $\langle x, x \rangle = 0, \langle x, y \rangle = 1, \langle y, y \rangle = 0$ or 1 .

We get orthogonal decomposition under \langle, \rangle :

$$H = H' \oplus (\mathbb{Z}x \oplus \mathbb{Z}y), \quad E = E' \oplus (\mathbb{Z}x \oplus \mathbb{Z}y)$$

Observe that $F \subset H'$ and $E \cap H' = E' = F^\perp \subset H'$, by the induction, we've finished our proof.

References

- [1] Browder, William, The Kervaire invariant of framed manifolds and its generalization. *Ann. of Math.* (2) 90 1969 157C186.
- [2] Browder, William, Surgery on simply-connected manifolds. *Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 65*. Springer-Verlag, New York-Heidelberg, 1972. ix+132 pp
- [3] Browder, William, Complete intersections and the Kervaire invariant. *Algebraic topology*, Aarhus 1978 (Proc. Sympos., Univ. Aarhus, Aarhus, 1978), pp. 88C108, *Lecture Notes in Math.*, 763, Springer, Berlin, 1979.
- [4] Buchstaber, Victor M.; Panov, Taras E. Torus actions and their applications in topology and combinatorics. *University Lecture Series*, 24. American Mathematical Society, Providence, RI, 2002. viii+144 pp. ISBN: 0-8218-3186-0
- [5] Fulton, William, Introduction to toric varieties. *Annals of Mathematics Studies*, 131. The William H. Roever Lectures in Geometry. Princeton University Press, Princeton, NJ, 1993. xii+157 pp. ISBN: 0-691-00049-2
- [6] Griffiths, Phillip; Harris, Joseph, Principles of algebraic geometry. Reprint of the 1978 original. *Wiley Classics Library*. John Wiley Sons, Inc., New York, 1994.
- [7] Kervaire, Michel A.; Milnor, John W. Groups of homotopy spheres. I. *Ann. of Math.* (2) 77 1963 504C537.
- [8] Kulkarni, Ravindra S.; Wood, John W. Topology of nonsingular complex hypersurfaces. *Adv. in Math.* 35 (1980), no. 3, 239C263.
- [9] Milnor, John W.; Stasheff, James D. Characteristic classes. *Annals of Mathematics Studies*, No. 76. Princeton University Press, Princeton, N. J.; University of Tokyo Press, Tokyo, 1974. vii+331 pp.
- [10] Ranicki, Andrew, Algebraic and geometric surgery. *Oxford Mathematical Monographs*. Oxford Science Publications. The Clarendon Press, Oxford University Press, Oxford, 2002. xii+373 pp. ISBN: 0-19-850924-3
- [11] Voisin, Claire, Hodge theory and complex algebraic geometry. I. Translated from the French by Leila Schneps. Reprint of the 2002 English edition. *Cambridge Studies in Advanced Mathematics*, 76. Cambridge University Press, Cambridge, 2007.
- [12] Wood, John W. Removing handles from non-singular algebraic hypersurfaces in $CP_{n+1}CP_n+1$. *Invent. Math.* 31 (1975), no. 1, 1C6.
- [13] Wood, J. W. Complete intersections as branched covers and the Kervaire invariant. *Math. Ann.* 240 (1979), no. 3, 223C230.
- [14] Libgober, A. S.; Wood, J. W. On the topological structure of even-dimensional complete intersections. *Trans. Amer. Math. Soc.* 267 (1981), no. 2, 637C660.
- [15] Wall, C. T. C. Classification of $(n-1)$ -connected $2n$ -manifolds. *Ann. of Math.* (2) 75 1962 163C189.

Wei WANG

School of Mathematical Sciences,

Fudan University,

Shanghai 200433, P.R. China.

Email: weiwang@amss.ac.cn